

# Simpson College International Student - Tuberculosis (TB) Screening Form

Name: \_\_\_\_\_ Birth Date: \_\_\_\_\_ Date Form Completed: \_\_\_\_\_

Please answer the following questions and return completed form to the address below, email, or fax provided.

Katie Lee, RN, BSN  
 Director of Student Health Services  
 Simpson College  
 701 North C Street  
 Indianola, Iowa 50125  
 USA

[katie.lee@simpson.edu](mailto:katie.lee@simpson.edu)  
 Fax: +1-515-961-1674  
 For questions call +1-515-961-1604

1. Have you ever had a positive TB skin test? Yes \_\_\_\_\_ No \_\_\_\_\_
2. Were you born in one of the countries listed below that have a high incidence of active TB disease? (If yes, please circle the country, below.) Yes \_\_\_\_\_ No \_\_\_\_\_
3. Have you had frequent or prolonged visits\*to one or more countries listed below with a high prevalence of TB disease? (If yes, please circle country.) Yes \_\_\_\_\_ No \_\_\_\_\_
4. Have you been a resident and/or employee of high-risk congregate settings (e.g. correctional facilities, long-term care facilities, and homeless shelters)? Yes \_\_\_\_\_ No \_\_\_\_\_
5. Have you been a volunteer or healthcare worker who served clients who are at increased risk for active TB disease? Yes \_\_\_\_\_ No \_\_\_\_\_
6. Have you ever been a member of any of the following groups that may have an increased incidence of latent M.tuberculosis infection or active TB disease: medically underserved, low-income, or abusing drugs or alcohol?  
 Yes \_\_\_\_\_ No \_\_\_\_\_

If the answer is **YES** to any of the above questions, Simpson College requires that you receive TB testing prior to your arrival on campus.

If the answer to all of the above questions is **NO**, no further testing or further action is required.

Afghanistan	Congo	Iran (Islamic Republic of)	Namibia	Singapore
Algeria	Côte d'Ivoire	Iraq	Nauru	Solomon Islands
Angola	Democratic People's Republic of Korea	Kazakhstan	Nepal	Somalia
Anguilla	Democratic Republic of the Congo	Kenya	Nicaragua	South Africa
Argentina	Djibouti	Kiribati	Niger	South Sudan
Armenia	Dominican Republic	Kuwait	Nigeria	Sri Lanka
Azerbaijan	Ecuador	Kyrgyzstan	Northern Mariana Islands	Sudan
Bangladesh	El Salvador	Lao People's Democratic Republic	Pakistan	Suriname
Belarus	Equatorial Guinea	Latvia	Palau	Swaziland
Belize	Eritrea	Lesotho	Panama	Tajikistan
Benin	Estonia	Liberia	Papua New Guinea	Thailand
Bhutan	Ethiopia	Libya	Paraguay	Timor-Leste
Bolivia (Plurinational State of)	Fiji	Lithuania	Peru	Togo
Bosnia and Herzegovina	French Polynesia	Madagascar	Philippines	Trinidad and Tobago
Botswana	Gabon	Malawi	Poland	Tunisia
Brazil	Gambia	Malaysia	Portugal	Turkmenistan
Brunei Darussalam	Georgia	Maldives	Qatar	Tuvalu
Bulgaria	Ghana	Mali	Republic of Korea	Uganda
Burkina Faso	Greenland	Marshall Islands	Republic of Moldova	Ukraine
Burundi	Guam	Mauritania	Romania	United Republic of Tanzania
Cabo Verde	Guatemala	Mauritius	Russian Federation	Uruguay
Cambodia	Guinea	Mexico	Rwanda	Uzbekistan
Cameroon	Guinea-Bissau	Micronesia (Federated States of)	Saint Vincent and the Grenadines	Vanuatu
Central African Republic	Guyana	Mongolia	Sao Tome and Principe	Venezuela (Bolivarian Republic of)
Chad	Haiti	Montenegro	Senegal	Viet Nam
China	Honduras	Morocco	Serbia	Yemen
China, Hong Kong SAR	India	Mozambique	Seychelles	Zambia
China, Macao SAR	Indonesia	Myanmar	Sierra Leone	Zimbabwe
Colombia				
Comoros				

Source: World Health Organization Global Health Observatory, Tuberculosis Incidence 2014. Countries with incidence rates of  $\geq 20$  cases per 100,000 population.

For future updates, refer to <http://www.who.int/tb/country/en/>

Prepared originally by ACHA's Tuberculosis Guidelines Task Force Revised April 2016 by Emerging Public Health Threats and Emergency Response Coalition

<http://www.acha.org/ACHA/Resources/Guidelines/ACHA/Resources/Guidelines.aspx?hkey=450d50ec-a623-47a2-aab0-5f011ca437fb>

\* The significance of the travel exposure should be discussed with a health care provider and evaluated.

Updated 04/2016